





# Know Diabetes by Heart™



Approximately 1 in 3 American adults with diabetes has CKD.



CVD risk is high in people with CKD.



Kidney disease is ranked in the top leading causes of death in the United States.



Every 24 hours, 170 people with diabetes begin treatment for kidney failure.

## Screening for Diabetes-Related Kidney Disease



### Who?

- ✓ Everyone with type 2 diabetes
- ✓ Everyone with type 1 diabetes for  $\geq 5$  years



### How?

UACR and eGFR



### When?

Annually


## Management of Diabetes-Related Kidney Disease

### Why Manage?

Decreases risk of CKD progression

Cardiovascular risk reduction

### Pillars of Therapy to Reduce Cardiorenal Risk



CKD = Chronic kidney disease  
 CVD = Cardiovascular disease  
 UACR = Urinary albumin to creatinine ratio  
 eGFR = Estimated glomerular filtration rate

RAAS Blockade = Renin-angiotensin-aldosterone system blockade  
 SGLT2-inhibitors = Sodium-glucose cotransporter 2 inhibitor  
 NS-MRAs = Nonsteroidal mineralocorticoid receptor antagonists  
 GLP-1 RAs = Glucagon-like peptide 1 agonists

Learn more at [KnowDiabetesbyHeart.org](https://www.knowdiabetesbyheart.org)