


Anxiety Disorders

7 A's Model



■ Be **AWARE** that people with diabetes may experience elevated anxiety symptoms

- › An anxiety disorder is a diagnosable mental condition characterized by frequent, intense, and excessive anxiety symptoms (for minimum six months).
- › Look for signs: frequent, intense, and excessive nervousness or worry, irritability, restlessness, trembling, dizziness, muscle tension, sleep disturbance, or panic attacks.

■ **ASK** about elevated anxiety symptoms

- › When to ask:
 - when the person reports symptoms or when you have noted signs (see **AWARE**)
 - in periods of significant diabetes-related challenge or adjustment (e.g., following diagnosis

of diabetes or complications, hospitalization, or severe hypoglycemia with loss of consciousness)

- during or after stressful life events
- if the individual has a history of anxiety disorder(s) or other mental health problems
- and in line with clinical practice guidelines.
- › Use open-ended questions or the Generalized Anxiety Disorder Two (GAD-2):
 - “Over the last two weeks, how often have you been bothered by feeling nervous, anxious, or on edge?”
 - “Over the last two weeks, how often have you been bothered by not being able to stop or control worrying?”
 - Response options are scored: “Not at all” (0), “Several days” (1), “More than half the days” (2), “Nearly every day” (3). Add the responses to form a total score.

- › If the GAD-2 total score is:
 - 3 or more—check whether the person has a current diagnosis of, or is currently receiving treatment for, an anxiety disorder. If neither, then further assessment is warranted (see **ASSESS**).
 - Less than 3 but you suspect a problem—consider whether the person may be experiencing diabetes distress, depression, or another psychological problem.

■ **ASSESS** for elevated anxiety symptoms using a validated questionnaire

- › When to assess:
 - in line with clinical practice guidelines
 - and if **AWARE** or **ASK** indicates elevated anxiety symptoms.
- › Use a brief, validated questionnaire, such as the Generalized Anxiety Disorder Seven (GAD-7).
- › GAD-7 total score 10 or more—moderate-to-severe anxiety symptoms are indicated:
 - inquire about any past history of anxiety disorder(s) and other mental health problems
 - consider the possibility of coexisting conditions (e.g., depression)
 - explore physiological, psychological, and behavioral causes for the elevated anxiety symptoms (including diabetes-related factors)
 - and continue to **ADVISE**.
- › GAD-7 total score less than 10—elevated anxiety symptoms are not indicated. Consider another psychological problem (e.g., diabetes distress, diabetes-specific fears, or depression).

NB: A clinical interview must be conducted (e.g., by a PCP or mental health professional) to diagnose an anxiety disorder—see **ASSIGN**.

NB: If you identify a person as having elevated anxiety symptoms, consider whether a suicide risk assessment is needed.

■ **ADVISE** about elevated anxiety symptoms

- › Explain that their questionnaire responses indicate they are experiencing elevated anxiety symptoms; they may have an anxiety disorder, which will need to be confirmed with a clinical interview.

- › Describe what an anxiety disorder is, and how it might affect their life overall and their diabetes management.
- › Advise that anxiety disorders are common, treatable, and can be managed effectively.
- › Advise that treating the anxiety disorder can help to improve their life overall and their diabetes management.
- › Make a joint plan about the “next steps” (e.g., what needs to be achieved and who can help).

■ **ASSIST** with developing an achievable action plan

- › If an anxiety disorder has been confirmed by clinical interview, assist the person to:
 - make an informed choice about a suitable treatment for the anxiety disorder—discuss the pros and cons for each option
 - set measurable goals to reduce anxiety symptoms and adapt their diabetes management plan if needed
 - identify sources of advice and support
 - start the preferred treatment (e.g., write a Mental Health Treatment Plan, a referral to a specialist for psychological support, or a prescription for medication).

■ **ASSIGN** to another health professional


- › If you do not feel comfortable or equipped to **ASSESS** or **ASSIST**, arrange a referral to a relevant health professional.
- › Maintain ongoing communication with the health professional to whom you made the referral.

■ **ARRANGE** follow-up care

- › Arrange a follow-up appointment to:
 - ask about the person’s progress
 - and assess the potential need for revising the action plan.
- › Be prepared to offer more support (e.g., through telephone reviews or more/extended consultations) during this time.

Depression

7 A's Model



■ Be **AWARE** that people with diabetes may experience depressive symptoms

- › Depression is a diagnosable mental condition characterized by a persistent state of low mood and lack of interest or pleasure in activities (for minimum two weeks).
- › Look for signs: lowered mood (e.g., sadness, hopelessness, or teariness), loss of interest or pleasure in usual activities, irritability, feelings of worthlessness or excessive/inappropriate guilt, difficulties concentrating, lack of energy, changes in weight and sleep patterns, psychomotor changes (e.g., moving or speaking more slowly than usual or being fidgety or restless), withdrawal from social and professional supports, reduced engagement in diabetes self-care tasks, and recurrent thoughts about death or suicide.

■ **ASK** about depressive symptoms

- › When to ask:
 - in line with clinical practice guidelines (e.g., on a routine or annual basis)
 - when the person reports symptoms or when you have noted signs (see **AWARE**)
 - in periods of significant diabetes-related challenge or adjustment (e.g., following diagnosis of diabetes or complications, hospitalization, or significant changes to the treatment regimen)
 - during or after stressful life events
 - or if the individual has a history of depression or other mental health problems.
- › Use open-ended questions or the Patient Health Questionnaire Two (PHQ-2):
 - *Over the last two weeks, how often have you been bothered by having little interest or pleasure in doing things?"*

- “Over the last two weeks, how often have you been bothered by feeling down, depressed, or hopeless?”
 - Response options are scored: “Not at all” (0), “Several days” (1), “More than half the days” (2), “Nearly every day” (3). Add the responses to form a total score.
- If the PHQ-2 total score is:
- 3 or more—check whether the person has a current diagnosis of, or is currently receiving treatment for, depression; if neither, then further assessment is warranted (see **ASSESS**).
 - Less than 3 but you suspect a problem—consider whether the person may be experiencing diabetes distress, elevated anxiety symptoms, or another psychological problem.

■ **ASSESS** for depressive symptoms using a validated questionnaire

- When to assess:
- in line with clinical practice guidelines (e.g., on a routine or annual basis)
 - or if **AWARE** or **ASK** indicates depressive symptoms.
- Use a brief, validated questionnaire, such as the Patient Health Questionnaire Nine (PHQ-9).
- PHQ-9 total score 10 or more—moderate-to-severe depressive symptoms are indicated:
- inquire about any past history of depression and other mental health problems
 - consider the possibility of coexisting conditions (e.g., anxiety disorder)
 - explore physiological, psychological, and behavioral causes for the depressive symptoms (including diabetes-related factors)
 - and continue to **ADVISE**.
- PHQ-9 total score less than 10—depressive symptoms are not indicated. Consider another psychological problem (e.g., diabetes distress or elevated anxiety symptoms).

NB: A clinical interview must be conducted (e.g., by a PCP or mental health professional) to diagnose major depression—see **ASSIGN**.

NB: It is essential that you conduct a suicide risk assessment if the person’s PHQ-9 total score is 10 or more, or they score 1 or more on item 9 of the PHQ-9.

■ **ADVISE** about depressive symptoms

- Explain that their questionnaire responses indicate they are experiencing depressive symptoms; they may have major depression, which will need to be confirmed with a clinical interview.
- Describe what depression is, and how it might impact on their life overall and their diabetes management.
- Advise that depression is common, treatable, and can be managed effectively.
- Advise that treating depression can help to improve their life overall and their diabetes management.
- Make a joint plan about the “next steps” (e.g., what needs to be achieved and who can help).

■ **ASSIST** with developing an achievable action plan

- If mild or major depression has been confirmed by clinical interview, assist the person to:
- make an informed choice about a suitable treatment for depression—discuss the pros and cons for each option
 - set measurable goals to reduce depressive symptoms and adapt their diabetes management plan if needed
 - identify sources of advice and support
 - and start the preferred treatment (e.g., write a Mental Health Treatment Plan, a referral to a specialist for psychological support, or a prescription for medication).

■ **ASSIGN** to another health professional


- If you do not feel comfortable or equipped to **ASSESS** or **ASSIST**, arrange a referral to a relevant health professional.
- Maintain ongoing communication with the health professional to whom you made the referral.

■ **ARRANGE** follow-up care

- Arrange a follow-up appointment to:
- ask about the person’s progress
 - and assess the potential need for revising the action plan.
- Be prepared to offer more support (e.g., through telephone reviews or more/extended consultations) during this time.

Diabetes Distress

7 A's Model



■ Be **AWARE** that people with diabetes may experience diabetes distress

- › Diabetes distress is the emotional burden arising from living with and managing diabetes.
- › Look for signs: suboptimal A1C or unstable blood glucose levels, missed clinic appointments, reduced engagement in diabetes self-care tasks, ineffective coping strategies, multiple life stressors, chronic stress, impaired relationships (personal or with health professionals), appearing passive/aggressive during consultations.

■ **ASK** about diabetes distress

- › When to ask: routinely ask all people with diabetes.
- › Use open-ended questions to explore the impact of diabetes on daily life and well-being, such as:
 - “What is the most difficult part of living with diabetes for you?”
 - “How is your diabetes getting in the way of other things in your life right now?”
- › If diabetes-related concerns are raised, continue to **ASSESS**.

■ **ASSESS** for diabetes distress using a validated questionnaire

- › When to assess:
 - annually, and
 - if **AWARE** or **ASK** indicates possible diabetes distress.
- › Use a validated questionnaire, such as the Problem Areas In Diabetes (PAID) scale or Diabetes Distress Scale (DDS)
 - an individual item score on either the PAID or DDS of 3 or more indicates a “problem area” or concern
 - a PAID total score of 40 or more out of 100, or a DDS score of more than 3.0 indicates severe diabetes distress.
- › If diabetes distress is not identified but a problem remains, consider whether another psychological problem is present (e.g., depression or anxiety).

■ **ADVISE** about diabetes distress

- › Explain diabetes distress.
- › Acknowledge that living with diabetes can be challenging; it is normal to have negative feelings about it, and to feel frustrated, anxious, distressed, or overwhelmed by diabetes.
- › Explain that learning about the signs and consequences of diabetes distress will help individuals recognize and respond promptly.
- › Make a joint plan about the “next steps” (e.g., what needs to be achieved and who can help).

■ **ASSIST** with developing an achievable action plan

- › Explore the individual’s source(s) of diabetes distress and the kind of support needed. For example, diabetes education, a review of their management plan, emotional/social/professional support, or a combination of these.

- › Provide relevant information and support.
- › Together, develop a plan for addressing their distress, prioritizing their most important or burdensome issues.
- › Consider the need for a referral if diabetes distress persists.

■ **ASSIGN** to another health professional


- › People with diabetes tend to prefer their diabetes health professional help them with the emotional impact of diabetes.
- › As diabetes distress is common and intertwined with diabetes management, it is best addressed by the diabetes health professional.
- › If you do not feel comfortable or equipped to do this, arrange referral to a relevant health professional to **ASSIST**.
- › Maintain ongoing communication with the health professional to whom you made the referral.

■ **ARRANGE** follow-up care

- › Arrange a follow-up appointment to:
 - ask about the person’s progress
 - and assess the potential need for revising the action plan.
- › If you used a questionnaire to **ASSESS** (e.g., PAID or DDS) consider using it again to reassess their level of diabetes distress. Agree on how often the reassessment would be helpful.
- › Be prepared to offer more support (e.g., through telephone reviews or more/extended appointments) during this time.

Eating Problems

7 A's Model



■ Be **AWARE** that people with diabetes may have eating problems

- › Eating problems occur on a continuum from full syndrome diagnosable eating disorders to disordered eating (behaviors).
- › Look for signs: frequent and restrictive dieting and problematic attitudes to food/eating, preoccupation and dissatisfaction with body weight/shape/size, unexplained weight gain or loss and other physical signs (e.g., calluses on hands, dental problems), less frequent (or no) self-monitoring of blood glucose, omission or restriction of insulin/medication, missed clinic appointments, recurrent diabetic ketoacidosis, erratic fluctuation of blood glucose levels, acute change of A1C (a sign of acute eating disorder onset and often insulin omission),

very high or very low A1C (in the case of food restriction while still using insulin), and concerns expressed by another person.

■ **ASK** about eating patterns and body image

- › When to ask: routinely ask all people with diabetes.
 - when the person raises a problem or concern, or when you have noted signs (see AWARE).
- › Use open-ended questions to talk about eating habits, weight, and body image, such as:
 - “Women [men] with diabetes are sometimes concerned about their weight or shape. How do you feel about your weight or shape?”
 - “Could you tell me a bit more about the recent changes in your eating patterns?”

- › Then, ask directly about insulin restriction/omission in a nonjudgmental way:
 - *“Some people with diabetes find it difficult to keep up with their insulin injections/boluses. How do you feel about it?”*
 - *“Some people miss or skip their insulin injections/boluses to manage their weight. Do you sometimes adjust your insulin to influence your weight?”* if so, *“How often does this happen?”*
- › Or use the mSCOFF.
- › If the person indicates a problem through open discussion, or responds “Yes” to one or more mSCOFF items, further assessment is warranted (see **ASSESS**).

■ **ASSESS** for eating problems

- › When to assess: if **AWARE** or **ASK** indicates a possible eating problem.
- › A comprehensive clinical assessment is required to diagnose the type and severity of the eating problem. This includes both a clinical interview and clinical examination. If this is outside of your expertise, you will need to refer the person to a health professional with expertise in eating disorders (see **ASSIGN**).

■ **ADVISE** about eating problems

- › Explain that the earlier conversation (or their mSCOFF score) indicates a potential eating problem, and further assessment is needed with a clinical interview.
- › Explain what disordered eating is and how this differs from an eating disorder.
- › Advise that untreated eating problems can negatively affect their life, but also their diabetes management and outcomes now and in the future.
- › Advise that eating problems are treatable, and can be managed effectively.
- › Advise that treating eating problems can help to improve their life overall and their diabetes management.

■ **ASSIST** with developing an achievable action plan

- › Provide information about the specific eating problem identified.
- › Explain treatment/therapy options.
- › Assist the person to access support and treatment.
- › Explain that a collaborative approach is needed, and who will be part of this multidisciplinary team and their role.
- › Agree on an action plan and achievable goals.
- › Make sure the person feels comfortable with the approach.

■ **ASSIGN** to another health professional


- › If the person is at immediate risk: refer them to hospital.
- › If the person is not at immediate risk, refer them to specialist eating disorder outpatient services or day programs if this is outside of your expertise.
- › Maintain ongoing communication with the health professional/service to whom you made the referral.

■ **ARRANGE** follow-up care

- › If you are part of the multidisciplinary team: continue to monitor the person’s progress (e.g., laboratory assessments and diabetes complications). Medical treatments, nutrition plans, and diabetes self-management goals will need to be adjusted regularly throughout the treatment.
- › If you are not part of the multidisciplinary team: inquire at each consultation about the person’s progress (e.g., have they engaged with the agreed treatment?).
- › Be prepared to offer more support (e.g., through telephone reviews or more/extended consultations) during this time.

Fear of Hypoglycemia

7 A's Model



■ Be **AWARE** that people with diabetes may experience extreme fear of hypoglycemia

- › Fear of hypoglycemia is specific and extreme fear evoked by the risk and/or experience of hypoglycemia.
- › Look for signs: “over-compensatory behaviors” (e.g., insulin dose reduction or overeating), “avoidance behaviors” (e.g., limiting exercise or social activities), an excessive number of daily blood glucose checks, acceptance of persistent high blood glucose levels, or not implementing “agreed” treatment changes to lower glucose levels.

■ **ASK** about experience and fear of hypoglycemia

- › When to ask:
 - about experience of hypoglycemia—at each appointment
 - about fear of hypoglycemia—when you have noted signs (see **AWARE**).
- › Use open-ended questions to talk about hypoglycemia and related fears, such as:
 - “What has been your worst experience with hypos?”
 - “What concerns you the most about hypos?”
- › Ask directly about compensatory behaviors:
 - “Some people take less insulin because they are worried about having a hypo. Do you ever reduce your insulin to avoid hypos?”
- › If the person indicates that they fear hypoglycemia, continue to **ASSESS**.

■ **ASSESS** for fear of hypoglycemia using a validated questionnaire

- When to assess: if **AWARE** or **ASK** indicates possible fear of hypoglycemia to further explore the specific concerns leading to fear.
- Use a validated questionnaire, such as the Hypoglycemia Fear Survey-II scale (HFS-II W).
 - For item scores 3 or 4, invite the person to explore their concerns. For example, *“I note here that you are concerned about [issue]. Can you tell me more about that?”* or, *“You seem to have a few worries about hypoglycemia. Which of these would you find most helpful to talk about today?”*
- Also assess whether the person purposefully keeps their blood glucose levels higher to reduce the risk of hypoglycemia.
- If fear of hypoglycemia is identified, continue to **ADVISE**.
- If fear of hypoglycemia is not identified, but a problem remains, consider whether another psychological problem is present (e.g., diabetes distress or anxiety).

■ **ADVISE** about fear of hypoglycemia

- Acknowledge the specific worries the person has raised under ASK or ASSESS.
- Acknowledge that it is common for people with diabetes to be concerned about hypoglycemia.
- Explain that a certain level of concern is adaptive and can motivate a person to act appropriately to treat or avoid hypoglycemia—but extreme fear needs attention.
- Advise that there are several ways to reduce fear of hypoglycemia. For example:
 - diabetes management strategies to reduce the frequency and severity of hypoglycemia
 - and psychological strategies that focus directly on extreme fear.
- If the person is persistently keeping their blood glucose levels in a higher range to avoid hypoglycemia, explain that this may have consequences for their long-term health.
- Make a joint plan about the “next steps” (e.g., what needs to be achieved and who can help).

■ **ASSIST** with developing an achievable action plan

- Through hypoglycemia/diabetes management:
 - review the person’s knowledge about hypoglycemia, how to treat, and barriers to treatment
 - review current diabetes management and discuss technologies (e.g., pump and continuous glucose monitor)
 - provide additional hypoglycemia training and/or diabetes education
 - and agree on an action plan to reduce hypoglycemia.
- Through fear management:
 - provide accurate information on their personal risk of hypoglycemia
 - provide support to change behaviors and unhelpful beliefs about hypoglycemia
 - and develop a stepwise plan to restore confidence and regain a sense of personal control.
- Consider the need for a referral if fear persists, or it is part of an anxiety disorder or the result of a traumatic hypoglycemia experience.

■ **ASSIGN** to another health professional


- People with diabetes tend to prefer their diabetes health professional help them with the emotional impact of diabetes.
- If you do not feel comfortable or equipped to do this, arrange referral to a relevant health professional to **ASSIST**.
- Maintain ongoing communication with the health professional to whom you made the referral.

■ **ARRANGE** follow-up care

- Arrange a follow-up appointment to:
 - ask about the person’s progress
 - and assess the potential need for revising the action plan.
- If you used a questionnaire to **ASSESS** (e.g., HFS-II W) consider using it again to reassess their level of fear of hypoglycemia. Agree on how often the reassessment would be helpful.
- Be prepared to offer more support (e.g., through telephone reviews or more/extended consultations) during this time.

Psychological Barriers to Insulin Use

7 A's Model



■ Be **AWARE** that people with type 2 diabetes may experience psychological barriers to insulin use

- Psychological barriers to insulin are the negative thoughts and feelings that people with diabetes may have about starting, using, or intensifying insulin.
- Look for signs: avoidance of talking about insulin (e.g., changing the topic, missing appointments), expressing concerns (e.g., about possible side effects, injecting, effect on lifestyle, perceptions of self and others), or insulin misuse (e.g., missing doses, smaller doses than recommended).
- Psychological barriers to insulin are experienced by people using insulin as well as those not yet using insulin.

■ **ASK** about psychological barriers to insulin use

- When to ask:
 - shortly after diagnosis of type 2 diabetes
 - when you notice signs of concerns or worries about insulin (see **AWARE**)
 - or if the person has suboptimal A1C despite being on (near) maximal oral agents.
- Use open-ended questions to discuss insulin and explore concerns, such as:
 - “How do you feel about going onto insulin [now or in the future]? Can you tell me more about that?”
 - “Some people have concerns or questions about insulin. Have you thought about insulin? Can you tell me more about that?”

- › If the person with type 2 diabetes indicates that they have questions or concerns about insulin, continue to **ASSESS**.

■ **ASSESS** for psychological barriers to insulin using a validated questionnaire

- › When to assess: if **ASK** indicates concerns or worries about insulin.
- › The Insulin Treatment Appraisal Scale (ITAS) is a validated measure. Barriers to insulin use are indicated by:
 - scores of 2 or less on positive appraisal items (3, 8, 17, 19)
 - or scores of 4 or more on negative appraisal items (the remaining items).
- › Invite the person to explore their concerns. For example, *“I note here that you are concerned about [issue]. Can you tell me more about that?”* or, *“You seem to have a few worries about insulin. Which of these would you find most helpful to talk about today?”*
- › If psychological barriers are raised, continue to **ADVISE**.

■ **ADVISE** about psychological barriers to insulin use

- › Acknowledge the specific barriers the person has raised under **ASK** or **ASSESS**.
- › Acknowledge that it is common to have questions and/or concerns.
- › Reassure them that needing insulin does not indicate they have “failed.”
- › Advise that many people need insulin as part of the natural progression of diabetes.
- › Tell them that people who use insulin find it beneficial because it:
 - is a powerful way to keep blood glucose within an optimal range to prevent long-term complications
 - allows for more flexibility in diet and planning of meals
 - and improves energy levels.

- › Advise that insulin use may begin with just one or two injections per day.
- › Make it clear that it is the individual’s decision whether or not to use insulin and you would like to assist them in making an informed choice.
- › Make a joint plan about the “next steps” (e.g., what needs to be achieved and who can help).

■ **ASSIST** with developing an achievable action plan

- › Assist the person with:
 - overcoming identified barriers to insulin using appropriate strategies (e.g., motivational interviewing techniques or an “insulin trial”)
 - making an informed choice about an action plan, including achievable and measurable goals for overcoming psychological barriers to insulin
 - and identifying sources of advice and support (e.g., structured group education or peer support).

■ **ASSIGN** to another health professional

- › People with diabetes tend to prefer their diabetes health professional help them with the emotional impact of diabetes.
- › If you do not feel comfortable or equipped to do this, arrange referral to a relevant health professional to **ASSIST**.
- › Maintain ongoing communication with the health professional to whom you made the referral.

■ **ARRANGE** follow-up care

- › Arrange a follow-up appointment to:
 - ask about the person’s progress
 - and assess the potential need for revising the action plan.
- › Be prepared to offer more support (e.g., through telephone reviews or more/extended appointments) during this time.